

# Wasabi - Japanese Horseradish

**G. P. Savage**

Food Group, Agriculture and life Sciences, Lincoln University, Canterbury, New Zealand.

## ABSTRACT

Wasabi, Japanese Horseradish (*Wasabia japonica* (Miq.) Matsum) is a perennial plant native to Japan. It has been cultivated in Japan for more than a thousand years and is now being grown in many countries as interest in Japanese cuisine expands. Wasabi can be grown in two main ways, either in flooded fields or in soil based mediums. The unique flavour of wasabi comes from isothiocyanates (ITCs) which are evolved from precursor glucosinolates by the enzyme myrosinase when the tissue is disrupted. ITCs found in wasabi are volatile, possess strong pungent smells and are toxic at high intakes. The overall flavour of wasabi depends on individual ITC content. Allyl ITC is found in the highest concentration in all tissues, ranging from 86-92% of the total ITC content. Apart from flavouring sauces and foods wasabi isothiocyanates have interesting anticancer effects. ITCs can also counter inflammatory conditions like asthma and anaphylaxis. ITCs have also been shown to inhibit platelet aggregation in the blood. Wasabi is a valuable crop that can be processed into a tasty condiment. Its production and consumption will increase as it becomes more appreciated in Western cuisine.

## INTRODUCTION

### History

Wasabi, known as Japanese Horseradish (*Wasabia japonica* (Miq.) Matsum) is a native condiment crop of Japan. It is not known when wasabi was first brought into cultivation but Japanese historical records indicate that wasabi, known originally as wild ginger, was introduced as a medicinal plant by Sukahito Fukae. The first Japanese medical encyclopaedia called "Honzo-wamyo" was published in A.D. 918 and it states that "wild ginger" (wasabi) had been grown in Japan for at least a thousand years (1). During 1596-1615 A.D. wasabi cultivation began on the upper reaches of the Abe River in Shizuoka prefecture. Its use, however, was reserved to the ruling class by order of the Shogun Ieyasu Tokugawa (2). At present, the natural distribution of wasabi in Japan ranges from Russia's Sakhalin island, north of Hokkaido (the most northern Japanese island) to Kyushu (the southernmost major Japanese island) (3). However, the Shimane region is the largest area of wasabi production and breeding research in Japan at present.

Wasabi is now being grown in many countries in the world including New Zealand, Taiwan, Korea, Israel, Brazil, Thailand, Columbia, near Vancouver, Canada and Oregon, USA. In New Zealand, the Ministry of Agriculture and Fisheries introduced wasabi for experimental cultivation in 1982 (4). Agronomic investigation of this crop was stimulated by commercial interest in 1986 (5,6). Preliminary assessment of the growth and plant yield of soil-grown wasabi was carried out at Lincoln in 1995 (7). Yields of flavour compounds, as affected by fertilizer treatment, were carried out in 1997 (8, 9). Further research has been performed at Lincoln University to develop an understanding of wasabi growing in New Zealand, especially the agronomy, cultivation methods, ITC variation and stability (10). Wasabi is now grown in New

Zealand for export of frozen rhizomes to Japan and it is also processed into a number of different wasabi flavoured food products.

### **Botany of wasabi**

Wasabi is a member of the *Cruciferae* family which also includes cabbage, cauliflower, broccoli, sprouts, water cress, radish, mustard and horseradish. The European horseradish (*Armoracia rusticana*) is known as a distant cousin of wasabi and is sometimes preferred by chefs, as a substitute for wasabi. The genus *Wasabia* consists of two species, *Wasabia tenuis* (an uncultivated species) and *Wasabia japonica* (the cultivated species). These two species are distinguished primarily by their cytology, stem size and colour and leaf size and shape (3). *Wasabia japonica* is a glabrous, perennial aromatic herb that grows about 450 mm high, producing leaves on long petioles from the crown of the plant. As the plant ages the rhizomes start to form and, at maturity after 18 months, the wasabi plant has a distinctive thickened stem (or rhizome) connected to the heart shaped leaves by long, thin petioles (Figure 1). Rhizomes are the most favoured plant part of wasabi. Most plants have one or two main rhizomes and a number of secondary stems (7). The lengths and weights of rhizomes vary widely between species e.g. for *Wasabia japonica* the length ranges from 50 to 200 mm long and weighs 4 to 120 g. *Wasabia japonica* leaves are simple, cordate-reniform, undulate-toothed and 80-250 mm in diameter. Petioles are vertical to oblique, 300–500 mm long, basally flattened and surround the rhizome. Whole fresh plants can weight up to 3.4 kg (3).

Wasabi flowers are white, bracteate, arranged on racemes, with ascending sepal, cruciform and obovate petals, perfect septum, elongate styles and simple stigma (3). Fertilisation is mainly by cross pollination. Seeds must be stored in a cool moist environment, since dry storage will result in desiccation and loss of viability of the seed. Fresh seed is naturally dormant and until it is vernalised by storing at a low temperature (5).

### **Wasabi cultivars**

In Japan a cultivar is usually named after its region of cultivation and, thus, wasabi cultivars are considered regionally specific in Japan. Seventeen wasabi cultivars have been developed and each has a strict cultivation and climate requirement that limits major cultivation to distinct areas (3). Specific regions and individual farmers produce their own unique cultivars as a result of persistent inbreeding and selection. According to some Japanese farmers wasabi has eight well-known cultivars which are Mazuma, Daruma, Takai, Shimane, Midori, Sanpoo, Izawa Daruma and Medeka. These cultivars were developed in the Shizuoka and Shimane prefectures. Another important cultivar ‘Hangen’ comes from the Kanagawa prefecture. Daruma is the most popular variety, known to grow well under marginal environmental conditions, such as warmer temperatures. It was developed by plant breeders based in the Shimane Research Station. For poor quality locations they developed Fuji Daruma, Izawa Daruma, Ozawa Daruma and Sanpoo in Shimane. However, all Shimane cultivars produce high quality rhizomes. Mazuma was developed in Shizuoka (but was originally grown in Wakayama and Okutama) and its production in Pacific Coast areas suggests that it has more heat tolerance than the Daruma cultivars although no published data can be found to verify this. In New Zealand, Daruma is the commercial cultivar of wasabi and preliminary studies using Shingen have just commenced.

### **Cultivation of wasabi**

In Japan, wasabi sometimes grows naturally in the gravel beds of mountain streams and is highly adapted to this environment. For commercial wasabi growing two types of cultivation methods are used. These are upland or soil grown wasabi and flooded field cultivation or water grown wasabi. Japanese growers select the method depending on the particular end use of the plants after harvest. Wasabi plants grown using the upland soil production method are harvested primarily for producing leaf and petiole products while wasabi plants grown using stream cultivation are harvested mainly for their enlarged rhizome, and only this plant part is used to make food products (3). Thus, it is believed in Japan that flooded systems can produce superior high quality enlarged stems. These are highly sought after and, therefore, command higher prices.

### **Soil grown wasabi**

Wasabi requires specific environmental conditions to thrive. Upland wasabi requires an air temperature from 6-20°C with 8-18°C considered optimal. Soils of pH 6-7 are considered best. It is most often grown on well-drained soil under mulberry or plum trees in Japan, whereas in New Zealand wasabi is usually grown in soil and in shade houses rather than under trees.

### **Water grown wasabi**

Stream grown wasabi requires air temperatures ranging from 8-18°C. However, a narrower range of temperatures (12-15°C) is considered ideal. An air temperature of less than 8°C inhibits plant growth and at less than 5°C plant growth ceases. Other factors can have an effect on the growth of wasabi and need to be considered carefully e.g. stable water temperature, good nutrient status in the water and well aerated, neutral or slightly acidic pH a high dissolved oxygen level and a large supply of water to maintain consistent flow (this particularly depends upon the field system being used). Rainfall accumulation is also important, with an even distribution desirable to stabilize the water supply and temperature (3). Spring water is considered best because of its clarity, stable temperatures and high level of oxygen. At warmer temperatures the dissolved oxygen in the water decreases, which inhibits the growth of plants. Silty or muddy water is undesirable as it may contain insufficient oxygen, but some silt in the water is considered beneficial as a source of nutrients. In Japan, wasabi grows on the wet banks of cool mountain streams and springs. Overall, construction and establishment of a flooded field is expensive and labour intensive. Water grown wasabi is produced in 42 prefectures, and soil grown wasabi in 21, out of 47 prefectures in Japan, which indicates that flooded cultivation is popular and is considered to produce a high quality product.

The unique environmental requirements and shortage of cultivatable lands limit wasabi production areas to 880 hectares in Japan (11) and 400 hectares in Taiwan, but demand for wasabi condiments is spreading from Japanese cuisine to modern western food. The increasing interest in wasabi and the inability to expand production in Japan has seen prices rise steadily since 1970 (3). High prices have stimulated research into soil production methods and the investigation of production areas outside Japan. Since 1982, the cultivation of wasabi has been trialled in New Zealand (6) because of New Zealand's climate (appropriate air temperature range, high quality water, long sunlight hours), which meets the ideal requirements for growing quality wasabi outside Japan. However, the area of soil grown wasabi is increasing not only in New Zealand but also in Taiwan, Colombia, Canada, Korea, Thailand and USA. The expansion in soil grown wasabi is mainly to reduce the high cost of flooded field establishment and associated high labour costs.

## Uses of wasabi

Wasabi adds a unique flavour, heat and greenish colour to foods and, thus, it is a highly valued plant in Japanese cuisine. Wasabi is described as having ‘a sharp hot taste with pungent smell’ but the heat component in wasabi is different from chillies, and the hotness quickly dissipates in the mouth leaving an extremely pleasant mild vegetable taste, with no burning sensation at all. Wasabi adds aesthetic and culinary appeal to many foods and is considered a staple condiment in the Japanese diet. Recently, it has found widespread appeal in western cuisine due to its ability to change an ordinary dish to an extra special one by improving the taste (with addition of a spicy flavour) and eye appeal i.e. by decorative contrast of the light green colour. As a result, it is becoming a new flavour for the rest of the world.

All the plant parts of wasabi possess some flavour but vary in the sharpness they deliver (8,12) and are, therefore, used for different purposes. Basically, wasabi can be served in three ways. These are as a condiment on the side of a dish, as a spice or herb in a dish and as wasabi flavour in processed foods. Rhizomes are the most popular tissues used to prepare fresh paste to be placed in a mound on a dish next to sliced raw fish (sashimi), spread on the raw fish in sushi preparations, or served on a small dish to accompany a bowl of cooked noodles (3). Sometimes grated wasabi is mixed with other ingredients like soya sauce and vinegar to prepare a dip for use with raw fish or other dishes, according to individuals’ choice. Tofu (soybean curd) is often decorated with grated wasabi.

Wasabi petioles and leaves are pickled in sake brine or soya sauce and are popular accompaniments for white rice. Sometimes fresh leaves are used in salads and dried leaves are used to flavour cheese, salad dressings or crackers. A wasabi wine is sold (mainly as a novelty) in some Japanese specialty stores as well as a high alcohol content wasabi liqueur.

Lower quality wasabi stems are commonly mixed with European horseradish (*Armoracia rusticana*) powder, mustard and food colour to produce ‘wasabi’ paste in tubes or to sell as wasabi powder. In New Zealand, a variety of genuine wasabi flavoured quality products e.g. sauces pasts and mayonnaise have been developed to add to snacks and foods. Genuine wasabi products contain only wasabi and are not diluted with European horseradish, beet extracts or mustard. If possible avoid wasabi products that are highly coloured with artificial additives (either bright green or yellow) as many of these products also contain other sources of taste which interfere with the clean taste of pure wasabi.

## Wasabi paste preparation

In traditional Japanese cuisine, wasabi is prepared by grating the fresh stem against a rough surface, such as a ginger grater, in much the same way as horseradish is prepared (3). But it is also noted that using sharkskin or “oroshi” as a tool for grating wasabi has been a practice in Japan since the earliest times and is still regarded as the preferred method of obtaining the best flavour, texture and consistency in freshly ground wasabi. Using a sharkskin grater and keeping the rhizome at a 90° angle to the grating surface is reported to minimize the volatiles’ exposure to the air. It is also stated that, in this way, the volatile compounds are allowed to develop with minimal dissipation. In New Zealand, wasabi paste is commercially prepared using mincers to finely grind the rhizomes and then it is mixed with other ingredients depending on the end use of the paste.

## Flavour constituents of wasabi

Isothiocyanates (ITCs) are a group of naturally occurring sulphur compounds responsible for the characteristic flavour of wasabi (13-15). The compounds are volatile in nature and are evolved from plant tissues when they are disrupted eg. in the preparation of food, grating, cutting, chewing etc. However, plant tissues do not contain ITCs, but contain glucosinolates which are the precursors of ITCs. Glucosinolates (GSL) are a group of glucosides (ie. they contain glucose in the structure), stored within the cell vacuoles of all *Crucifereae* plants (14). Glucosinolates are a complex group of  $\beta$ -D thioglucose compounds synthesised from amino acids (8). They contain a sulphonic group which is usually bound to sodium or potassium making them anionic (16, 17, 18). The sulphate in the sulphonic group is attached through a C=N bond and different side groups (R) give a wide range of related glucosinolates. Each one has its own characteristic odour or taste (Table 1). Due to the presence of the glucose in the molecule glucosinolates are hydrophilic, non volatile compounds (17).

When plant tissues are mechanically disrupted or injured (eg. by chewing, crushing or grating in the preparation of food), the myrosinase is released from the cell wall and in the presence of adequate moisture myrosinase rapidly hydrolyses the GSLs (Figure 2) to yield glucose and an aglucone. Some of the intermediate steps have not been fully described (19, 20). The organic aglucone is unstable and undergoes Lossen Rearrangement (21) to produce sulphate and a variety of other products.

The nature of the products is dependent on the number of factors, including the structure of the GSL side chain, the reaction conditions (eg. pH), the presence of cofactors (eg. metal ions, specific proteins), temperature and duration as well as the age and condition of the plant tissues. Isothiocyanates (ITCs) are formed from GSLs under neutral and alkaline conditions. However, GSLs that contain a  $\beta$ -hydroxyl group (I in Figure 2) in their side chain, give rise to ITCs that spontaneously cyclize to form oxazolidinethiones. Some aromatic and heterocyclic GSLs (II in Figure 2) produce ITCs which are unstable at pH 7 or higher and break down to release the corresponding alcohol and inorganic thiocyanate ions.

However, once formed, ITCs are more stable under acidic conditions. In weakly acidic pH or in the presence of  $\text{Fe}^{+2}$  and/or endogenous nitrile factor, nitriles are produced from aglucone by autolysis instead of ITC, with the liberation of elemental sulphur (20). The relative proportion of ITC to nitriles can vary widely depending upon the conditions of autolysis (20). Thiocyanate formation is believed to involve a cofactor, which may also be a protein, since it has been shown to be labile to both heat and polar organic solvents. Most of the sulphur containing end products formed by the enzymatic and non-enzymatic reactions of GSLs are volatile (14).

Several ITCs have been reported from previous investigations into wasabi and each ITC has a specific flavour profile (8, 15) with the complete taste of wasabi being derived from the combined tastes and odours of all the ITCs present. A summary of different ITCs reported from previous investigations into the ITC content of wasabi tissue is listed in Table 1

Allyl ITC has the main effect on the overall taste of wasabi because it is the ITC found in highest concentration in the rhizomes and other plant tissues (9). Allyl ITC is also found in the highest concentration in horseradish (12). While allyl ITC is the main flavour component of wasabi due to its pungency other ITCs e.g. 6-methylthiohexyl ITC and 7-methylthioheptyl ITC,

by giving their characteristic fresh greenish flavour, may contribute significantly to the total taste profile of wasabi (15, 24).

### **Medicinal properties of isothiocyanates**

The medicinal value of chemicals extracted from wasabi were first documented in the Japanese medicinal encyclopaedia during the 10<sup>th</sup> century (24). Recently, research interest in ITCs has become more intense because of their potential to have a wide variety of medicinal, pharmacological or industrial applications. These exciting applications are at an early stage of investigation, most likely because of wasabi's high present commercial value and scarcity.

### **Anticancer effects**

Medicinally, the most important feature of ITCs is evidence that they have a chemopreventive effect on cancer at a variety of organ sites including lung, mammary glands, liver, oesophagus, bladder, pancreas, colon and prostate (25). A recent case-control study in Los Angeles (26) showed that high consumption of *cruciferae* vegetables containing ITCs reduced the risk of developing colon cancer (27-31). Tests have been carried out on tumours in rats and it has been reported that some ITCs have a protective role against breast, stomach and colon cancers in rats (32, 33). Several mechanisms have been proposed and investigated for tumour inhibition by ITCs, for instance sulfoforane (SFN) and phenylethyl ITC (PEITC) are reported as potent inducers of the phase II enzymes involved in detoxification of carcinogens (34). The inhibition of chemically induced lung tumorigenesis by PEITC was mediated primarily by the inhibition of metabolism which resulted in a decrease in O<sup>6</sup> methylguanine in lung DNA, indicating that ITC targets cytochrome P450s (35, 36). Results from recent bioassays in A/J mice appear to support the mechanism of induction of apoptosis in lung by PEITC and butyl ITC (BITC) (37). Extensive research on ITCs commonly found in cruciferous vegetables such as watercress, broccoli, radish and cabbage have been linked to the reduced risk of certain human cancers (27, 28). From the chemoprevention point of view it is important to know whether the beneficial effects come, at least in part, from ITCs in the diet. In a cohort study it was clearly shown that individuals with detectable levels of ITCs in the urine were less likely to develop lung cancer (38). It has been suggested that normal dietary levels of ITCs derived from wasabi or other fresh cruciferous vegetables eaten regularly can protect against the low levels of carcinogens encountered in everyday life (25). As a result, the American Cancer Society recommends that cruciferous vegetables should be part of every person's daily diet to reduce the risk of several cancers.

### **Effects of isothiocyanates on blood**

The ITCs in wasabi have been tested for inhibition of platelet aggregation mediated by arachidonic acid (39), and for deaggregation. ITCs showed a ten times higher response than is reported for aspirin. In the case of heart attacks, where aspirin is commonly prescribed, ITCs have been shown to have a more rapid action at low levels than the thirty minutes for aspirin. In this regard, the most potent ITCs reported are  $\omega$ -methylthioalkyl ITCs, especially 6-methylthiohexyl followed by 7-methylthioheptyl and 5-methylthiopentyl ITCs. The anticoagulant property of ITCs could be used in the treatment of elderly people and during surgery where preventing platelet aggregation is vital for the well being of the patient. The mechanism by which the ITCs inhibit platelet aggregation from occurring has not been precisely

determined but may be through a specific inhibition of the arachidonic acid cascade (40). This therefore raises the possibility of using ITCs to limit inflammation in tissues.

### **Anti-asthmatic and anti-inflammatory properties**

Benzyl and allyl ITCs from onion extracts showed anti-asthmatic effects when studied by Dorsch *et al.* (41). Thromboxanes (generated by lung tissue and by aggregating platelets during lung anaphylaxis) and prostaglandins (generated by mast cells during activation) are known to cause bronchial obstruction and generally play a role in the pathogenesis of bronchial asthma. The isothiocyanates prevented bronchial obstruction caused by subsequent inhalation of ovalbumin but did not prevent obstruction caused by inhalation of histamine acetylcholine. This indicates that the anti-asthmatic effect of ITCs are not due to an anti-histamine effect but act by inhibiting the inflammatory process at an earlier stage, possibly the production or action of other inflammatory molecules such as thromboxanes or prostaglandins. Thus, ITCs could potentially be used to counter inflammatory conditions such as asthma or even anaphylaxis. Traditionally, horseradish and mustard have been used as a remedy for clogged sinuses, relief of congestion, muscular pain and inflamed joints (40).

### **Antibiotic effect**

Masuda (42) has suggested that wasabi could contribute to a healthy smile by inhibiting the growth of the bacteria on teeth and gums in the mouth. *Streptococcus mutans* is known to cause dental caries and the consumption of wasabi can reduce bacterial activity. This was explained by wasabi's ability to interfere with the sucrose-dependent adherence of cells to the surface of teeth and gums.

### **Other industrial applications**

ITCs extracted from wasabi can be used to make antibiotics, fungicides, insecticides, nematocides and as wood preservatives (43). ITCs are said to act as an antidote to food poisoning bacteria, one factor that has led to the use of wasabi with raw fish dishes in Japan (44-46). It is also reported that ITCs may have a role in protecting against diarrhoea (47). ITCs have also been used as antifouling compounds to stop seaweed growing on ships' hulls. Recently it has been shown that wasabi contains anti-fungal metabolites that can render plants resistant to virulent isolates of the blackleg fungus (48). This fungus can devastate commercially important crops such as the oilseed plants rapeseed and canola. There is a potential to develop a natural fungicide using wasabi extracts.

### **Conclusions**

Wasabi is a valuable crop that can be made into a tasty condiment. In Japanese cuisine freshly ground wasabi is used to add a spicy flavour directly to foods. Wasabi is also made into a paste which is stabilised by the addition of a number of other ingredients. In Western cuisine where hot spicy tastes are a recent addition to the diet, milder sauces and mayonnaises containing wasabi are more often appreciated. While the main reason for consuming wasabi is the unique taste, it is interesting to note that the active components, the isothiocyanates appear to have some positive anti-cancer effects in the body. This will lead to increased production and consumption of this interesting perennial crop.

### **Acknowledgements**

Thanks to Tamanna Sultana for her assistance in the preparation of this paper.

## REFERENCES

- 1- Hodge WH. 1974. *Econ Bot* 28: 118-129.
- 2- Kojima M. 1981. *Nogyo Oyobi Engei*, 56(5): 113-118.
- 3- Chadwick CI, Lumpkin TA and Elbersen LR. 1993. *Econ Bot* 47: 113-135.
- 4- Forde B. 1982. Department of scientific and Industrial research discussion paper no. 7. Wellington, NZ, 25p.
- 5- Palmer J .1990. *N Z J Crop Hort Sci* 18: 161-164.
- 6- Douglas J.A. and Follett JM. 1992. *Proc Agron Soc* 22: 57-60.
- 7- Martin RJ and Deo B. 2000. *N Z J Crop Hort Sci* 28: 45-51.
- 8- Sultana T, Savage GP, McNeil DL, Porter N and Martin RJ. 2000. *Proc Nutr Soc NZ* 25: 95-106.
- 9- Sultana T, Savage GP, McNeil, DL, Porter, N, Martin RJ and Deo B. 2002. *J Sci Food Agric* 82:1477-1482.
- 10-Sultana T, Porter NG, Savage GP McNeil DL 2003. *J Agrc Food Chem* 51: 3586-3591.
- 11- Adachi S. 1987. Wasabi cultivation. Shujyunsha Co. LTd., Tokyo, Japan. P 199. (Translated).
- 12- Sultana T, Savage GP, McNeil DL, Porter N and Clark B. 2002. *Inter J Food Sci Nutr* (Submitted).
- 13- McGregor DI, Mullin WJ and Fenwick GR 1983. *J Ass off Anal Chem* 66 : 825-849.
- 14- Delaquis PJ, Mazza G. 1995. *Food Technol* 73-84.
- 15- Masuda H, Harada Y, Tanaka K, Nakajima M, Tebeta H, Takeoka GR, Teranishi R, Williams PJ and Kobayash IA 1996. *Biotech improved foods flavours*. ACS Symposium Series No. 637 25: 67-78.
- 16- Pretera T, Fahey JW, Holtzclaw WD, Abeygunawardana C, Kalchinski JL and Talalay P. 1996. *Anal Biochem* 239: 168-179.
- 17- Larsen PO. 1981. Glucosinolates in the biochemistry of Plants. A comprehensive Treatise, Stumpf PK and Conn, EE Eds, Vol 5, pp 502-525, Academic press, N Y.
- 18- Warton B, Matthiessen JN, Shackleton MA. 2001. *J Agric Food Chem* 49: 5244-5250.
- 19- Fahey JW, Zalcmann AT and Talalay P. 2001. *Phytochem* 56: 5-51.
- 20- McGregor DI 1993. In: Encyclopaedia of Food Science Food Technology and Nutrition, Macrae R, Robinson RK and Sadler MJ. Eds, Vol 4, Academic press, Harcourt Brace Jovanovich publishers.
- 21- Fenwick R, Heaney RK, Mullin WJ. 1983. *CRC Crit Rev Food Sci Nutr* 18: 123-201.
- 22- Kumagai H, Kashima N, Seki T, Sakurai H, Ishii K and Ariga T. 1994. *Biosci Biotech Biochem* 58 (12): 2131-2135.
- 23- Etoh H, Nishimura A, Takasawa R, Yagi A, Satio K, Sakata K, Kishima I and Ina K. 1990. *Agric Biol Chem* 54: 1578-1589.
- 24- Ina K, Sano A, Nobukuni M and Kishima I. 1981. *Nippon Shokuhin Kogyo Gakkaishi* 28: 365-370.
- 25- Chung FL. 2002. Chapter 7 in *Phytochemicals in Nutrition and Health* edited by Meskin MS, Bidlack WR, Davies AJ and Omaye ST, CRC Press USA.



- 26- Lin HJ, Probst-Hensch NM, Louie AD, Kau IH, Witte JS, Ingles SA, Frankl HD, Lee ER and Haile RW. (1998). *Cancer Epid Biomark Prev* 7: 647-652.
- 27- Verhoeven DTH, Goldbohm RA, van Poppel G, Verhagen H and van den Brandt PA. 1996. *Cancer Epid Biomark Prev* 5: 733-748.
- 28- Steinmetz KA and Potter JD. 1991. Vegetables, fruit, and cancer. II: Mechanisms. *Cancer Causes Control* 2: 427-442.
- 29- Tanida N, Kawaura A, Takahashi A, Sawada K and Shimoyama T. 1991. *Nutr Cancer* 16: 53-58.
- 30- Fuke Y, Haga Y, Ono H, Nomura T and Ryoyama K. 1997. *Cytotech* 25: 197-203.
- 31- Fuke Y, Ohishi Y, Iwashita K, Ono H and Shinohara K. 1994. *J Japanese Soc Food Sci Technol* 41:709-711 (In Japanese with English summary).
- 32- Wattenberg LW. 1977. *J Nat Cancer Inst* 58: 359-398.
- 33- Wattenburg LW. 1981. *Cancer Res* 41: 2991-2994.
- 34- Zhang Y and Talalay P. 1994. *Cancer Res* 9 (suppl.), 54: 1976-1981.
- 35- Morse MA, Amin SG, Hecht SS and Chung FL. 1989. *Cancer Res* 49: 2894-2897.
- 36- Morse MA, Eklind KL, Hecht SS, Jordan KG, Choi CI, Desai DH, Amin SG and Chung FL. 1991. *Cancer Res* 51: 1846-1850.
- 37- Yang YM, Conaway CC, Wang CX, Dai W, Chiao JW, Albino AP and Chung FL. 2002. *Cancer Res*, In press.
- 38- Seow A, Shi C-Y, Jiao D, Hankin JH, Lee H-P, Coetzee GA and Yu M. 1998. *Cancer Epid Biomark Prev* 7:775-781.
- 39- Kumagai H, Kishima N, Seki T, Sakurai, H., Ishii K and Ariga T. 1994. *Biosci Biotechnol Biochem* 58: 2131-2135.
- 40- Depree JA Howard TM and Savage GP. 1999. *Food Res Inter* 31: 329-337.
- 41- Dorsch W, Adam O, Weber J. and Ziegeltrum T. 1985. *Euro J Pharmacol* 107: 17-25.
- 42- Matsuda H. 2000. *Sci News* 159: 29.
- 43- Brown PD and Morra MJ. 1997. *Adv Agron* 61: 167-231.
- 44- Ono H, Tesaki S, Tanabe S and Watanabe M. 1998. *Biosci Biotech Biochem* 62(2): 363-365.
- 45- Shin IS and Lee JM. 1998. *Bull Korean Fish Soc* 31: 835-841.
- 46- Hasegawa N, Matsumoto Y, Hoshino A and Iwashita K. 1999. *Internat J Food Microbiol* 49: 27-34.
- 47- Nakayama H, Suzuki T and Suzuki Y. 1998. *Nippon Nogeikagaku Kaishi* 72: 499-507.
- 48- Soledade M, Pedras C and Sorensen JL. 1998. *Phytochem* 49: 1959-1965.